
Rings Node Report

Mark Showalter & Mitch Gordon PDS Management Council Meeting Washington, DC March 25, 2010

Cassini CIRS Re-formatting

- Pipeline rebuilt to satisfy liens.
- New diagrams generated.
- Entire data set regenerated from COCIRS 2.0.
- The review panel missed a few things...
 - E.g., each spectrum now includes an indication of whether the shutter was open or not!



Cassini CIRS Re-formatting

Lessons Learned

• Get the team to do it right the first time.


Cassini CIRS Re-formatting

Lessons Learned

- Get the team to do it right the first time.
 - This is way easier said than done.


Cassini UVIS Occultations

Supported by CDAP funding to Josh Colwell



Uncalibrated:
Photon counts vs. time,
binary formats


Calibrated:
Optical depth vs. radius,
ASCII formats



- Our Original Goals:
 - Fully integrate HST data into OPUS.
 - Mainly images of outer planets.
 - Let users download HST data directly from an OPUS search.
 - Produce usable, PDS-compliant, complete metadata and indices.

- The Problem:
 - STScl revised every data product before we finished.
 - STScl updates their calibration pipeline and file formats whenever they like, without warning.

- Our Original Revised Goals:
 - Fully Partially integrate HST data into OPUS.
 - Let users download preview HST data directly from an OPUS search.
 - ... and provide instructions on how to request the data from STScl.
 - Produce usable, PDS-compliant nearly complete metadata and indices.




```
****************************
fitsfile= 'J9DU03B3Q DRZ.FITS' / name of file from the STScI archive
headerno=
                          1 / header number in file, starting at 1
recordno=
                           1 / record number in file, starting at 1
                            T / conforms to FITS standard
SIMPLE =
                           8 / array data type
BITPIX =
NAXIS =
                           0 / number of array dimensions
EXTEND =
ORIGIN = 'NOAO-IRAF FITS Image Kernel July 2003' / FITS file originator
IRAF-TLM= '21:12:50 (25/09/2009)' / Time of last modification
NEXTEND =
                            3 / Number of standard extensions
     = '2009-09-25T21:12:50' / date this file was written (yyyy-mm-dd)
DATE
FILENAME= 'j9du03b3q drz.fits' / name of file
FILETYPE= 'SCI
                           / type of data found in data file
TELESCOP= 'HST'
                             / telescope used to acquire data
INSTRUME= 'ACS '
                             / identifier for instrument used to acquire data
EQUINOX =
                       2000.0 / equinox of celestial coord. system
             / DATA DESCRIPTION KEYWORDS
ROOTNAME= 'j9du03b3g'
                             / rootname of the observation set
IMAGETYP= 'EXT
                            ' / type of exposure identifier
PRIMESI = 'ACS '
                           / instrument designated as prime
             / TARGET INFORMATION
TARGNAME= 'JUPITER3
                                       ' / proposer's target name
RA TARG = 2.259050923144E+02 / right ascension of the target (deg) (J2000)
DEC TARG= -1.592159661850E+01 / declination of the target (deg) (J2000)
             / PROPOSAL INFORMATION
PROPOSID=
                        10507 / PEP proposal identifier
LINENUM = '03.001
                       ' / proposal logsheet line number
                                      ' / last name of principal investigator
PR INV L= 'Grodent
PR INV F= 'Denis
                              ' / first name of principal investigator
                              ' / middle name / initial of principal investigat
PR INV M= '
             / EXPOSURE INFORMATION
SUNANGLE=
                   138.593323 / angle between sun and V1 axis
MOONANGL=
                   109.851707 / angle between moon and V1 axis
                   -16.545755 / altitude of the sun above Earth's limb
SUN ALT =
FGSLOCK = 'FINE
                            ' / commanded FGS lock (FINE, COARSE, GYROS, UNKNOWN)
                             / number of gyros scheduled for observation
GYROMODE= '2'
REFFRAME= 'GSC1
                            / quide star catalog version
DATE-OBS= '2006-03-27'
                            / UT date of start of observation (yyyy-mm-dd)
TIME-OBS= '09:01:53'
                             / UT time of start of observation (hh:mm:ss)
               53821.37630852 / exposure start time (Modified Julian Date)
EXPSTART=
               53821.37753111 / exposure end time (Modified Julian Date)
EXPEND =
                        105.0 / exposure duration (seconds) -- calculated
EXPTIME =
EXPFLAG = 'NORMAL
                             / Exposure interruption indicator
```


- Our Solution:
 - Every HST observation is now a DOCUMENT object.
 - The document includes TIFFs and JPEGs of the images (but not the FITS files).
 - The document contains all the FITS header parameters from the files.

OPUS Development


"Outer Planets Unified Search"



- We remain current with Cassini deliveries
- Most recent changes are "under the hood"
 - Performance issues
 - REST-ful interface




On every user click...




On every user click...

The result count updates.




On every user click...

- The result count updates.
- The tab selection updates.



On every user click...

- The result count updates.
- The tab selection updates.
- Remaining options that would return no hits turn gray.




"Asynchronous Communication"

The advantage:

- The user gets immediate feedback about available data when exploring.
- Users like it.


The disadvantage:

 Our server has to handle potentially dozens of queries on the database all at once.



REST-ful URLs

```
http://pds-rings.seti.org
/search/
 #planet =
      Saturn
 &target =
      EPIMETHEUS
 &instrumentid =
      Cassini+ISS
&tab =
      obs instrument
        COISS
 &open =
      planet, target,
      instrumentid,
       FILTER, camera
```





REST-ful URLs

The advantage:


- This begins the decoupling of the user interface from the search <u>service</u>.
- You can bookmark any step in a search.
- ANYBODY can create a web-based tool that accesses our service.

Geometric Metadata






...based on a fine sampling of the entire field of view



Web Site Re-design...



SETI Institu

+ Carl Sagan Center for the Study of Life in the Universe

Node Manager: Mark R. Showalter Webmaster: Neil Heather PDS Atmospheres

Geosciences

Imaging Na

Navigation

Plasma/Particle Interactions

Rings Small Bodies

SETI INSTITUTE

SETI Institute

+ The Carl Sagan Center for the Study of Life in the Universe

Node Manager: Mark R. Showalter Webmaster: Neil Heather NASA Official: William P. Knopf

ywords for Rings Node internal indexing: planetary rings, planetary rings data, planetary rings images, planetary rings spectra, jupiter rings, jupiter rings data, jup gs images, jupiter rings spectra, saturn rings, saturn rings data, saturn rings images, saturn rings spectra, uranian rings, uranian rings data, uranian rings images • NASA Portal

• Site Help

• Feedback

Planetary Rings Node

Home | About Us | About PDS | About SETI | News | Education | Data & Information | Research Support | Contact Us

OPUS: Outer Planets Unified Search

Quick Start Page


Click any relevant boxes, watch the observation count at right, then click "Go!" to see results.

Target Requirements:	Data Source:	Measurement Needs:	Observa S
Planet:	From mission:	Measurement quantity:	Break d
✓ Jupiter	☐ Cassini	Intensity or reflectivity	Target
Saturn	Earth-based Telescope	Opacity	Adra
Uranus	✓ Galileo	Spatial sampling type:	Ama
	☐ Hubble Telescope☐ New Horizons	Profiles (1-D)	
Target type:	☐ Voyager	Images (2-D)	lo
Planet	□ voyagei	Extended 2-D	Euro
Ring	Instrument type:	("mosaics")	Gar
Moon	✓ Camera	Wavelength band:	Call
Star	□ Near-IR Spectrometer	☐ Visual	
☐ Sky	☐ Thermal IR	☐ Near IR	Instrun
Other	Spectrometer	☐ Thermal IR	Gali
or enter a target name:	UV Spectrometer	Radio	
	High-Speed	□UV	Select.
	Photometer	Wavelength sampling type:	
or select target from list	or select mission &	Detailed ("spectra")	
3	instrument name from <u>list</u>	Limited ("color")	
would also like to limit my search further based on Subregion on the target and/or spatial resolution Lighting and viewing geometry on the target Wavelength and bandpass Time limits and duration Mission- and instrument-specific parameters Refine query further		Timespan: Minutes ("movies") Days Years Time sampling: Minutes Hours Days	iono Di

OPUS Help Update & change history Submit a comment/report a bug

Result Summary			
Observation count: 3202 Show me my results			
Break down results by			
Target name Adrastea	5		
Adrastea Amalthea	5 4		
lo	104		
Europa	1202		
Ganymede	877		
Callisto	1010		
Instrument	▼		
Galileo SSI	3202		
Select	▼		

Small Bodies



Web Site "Facelift"

- Goals: Ease of use, ease of maintenance, consistency, user appeal, compatibility with other nodes
- Standardized page templates and a Content Management System

Node Internal Matters

- We're adopting Python/Django in a big way.
- Compared to equivalent C or FORTRAN, many codes are ~ 10 x shorter.
 - There's a library for everything!
- In-house development includes a CSPICE interface and a PDS3 label parser.
- It's FREE.
 - cf. IDL license fees.

Coming Soon...

SETI Institute will be relocating this summer.